

Substance Abuse in Brief Fact Sheet

Fall 2006, Volume 4, Issue 2

Identifying and Helping Patients With Co-Occurring Substance Use and Mental Disorders: A Guide for Primary Care Providers

It is estimated that approximately 4.6 million American adults have both substance use disorders (SUDs) and mental disorders (MDs).¹ Other terms for these co-occurring disorders (CODs) are dual diagnosis and comorbidity. Early recognition and treatment of these CODs are essential to improving treatment outcomes and quality of life for these patients.² This publication will help primary care providers identify patients with CODs and provide appropriate treatment for these patients.

Identifying Patients With COD

Primary care providers, such as physicians, physician's assistants, and nurse practitioners, are in an excellent position to help their patients with both SUDs and MDs. These disorders may exacerbate or be related to other health problems, such as headaches, cardiovascular disease, high blood pressure, diabetes, digestive disorders, and cirrhosis,^{3,4} so patients with CODs may often seek medical care from primary care providers. Hence, primary care providers may have established a relationship with their patients conducive to discussing SUDs and MDs.

Barriers to identifying and treating co-occurring disorders in the primary care setting include the following:

COD Red Flags^{11, 13, 15, 16}

- Nasal irritation (SUD only)
- Unexplained bruises (SUD only)
- Enlarged liver or spleen, abnormal liver function, hepatitis, or cirrhosis in later stages (SUD only)
- Withdrawal symptoms (SUD only)
- Headaches
- Chest pain or cardiac arrhythmia
- Gastrointestinal symptoms
- Hypertension
- Sexual dysfunction
- Fatigue
- Apathy or flat affect
- Social withdrawal
- Changes in concentration, mood, activity level, sleeping, appetite, or weight
- Feeling of worthlessness or inappropriate guilt
- Fear, worry, or repetitive, intrusive thoughts or actions
- Problems with cognition or impulse control
- History of physical or mental trauma

*These symptoms appear in combination and may indicate other problems as well as COD.

- **Low rates of screening**,^{5, 6} **correct diagnosis**,^{7, 8} and **appropriate referral**.⁵
- **Patient denial**.⁹ Patients may not realize they have a problem. Approximately one-third of patients do not disclose self-perceived SUDs or MDs to their primary care provider.¹⁰
- **Patient reluctance to talk to the provider**. The stigma of CODs still exists, and many patients are afraid to admit they have a problem.³
- **Provider attitudes**.^{7, 11} Providers may fail to make the possible link between physical symptoms and SUDs and/or MDs when assessing a physical problem. Providers' preconceptions and attitudes can also hinder identification of such disorders.
- **Inadequate training**. Many providers receive inadequate education on CODs and their effects on other medical conditions.³
- **Time constraints due to short appointment times**.⁹

Some barriers can be overcome by incorporating three components into the primary care practice: (1) obtaining an annual patient history on substance use and mental health issues, (2) being aware of warning signs that may be related to an SUD or MD, and (3) screening routinely for CODs.

Patient History

All patients should complete an annual health history, including questions about personal and family history of substance use and mental health issues.¹² Questions about victimization, trauma, personal, and social issues (e.g., unemployment, legal problems, homelessness, financial or marital difficulties) should be included because these can be related to SUDs and MDs.^{11, 12, 13} Primary care providers should ask questions about substance use and mental health symptoms, preferably in the context of other lifestyle questions so that these potentially sensitive topics seem less threatening to patients.¹⁴ An open, empathetic, and nonjudgmental attitude is essential to encouraging patients to talk about their symptoms.¹¹



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Substance Abuse and Mental Health Services Administration
Center for Substance Abuse Treatment
www.samhsa.gov

Substance Abuse in Brief Fact Sheet

Fall 2006, Volume 4, Issue 2

Warning Signs

Most patients will not visit a primary care setting with obvious, immediate signs of an MD, such as delirium, confusion, or disorientation, or an SUD, such as odor of alcohol on the breath or marijuana on clothing, dilated pupils, slurred speech, or needle marks. However, warning signs—“red flags”—for SUDs and MDs can manifest in subtle physical or behavioral symptoms (see sidebar on front page). For example, many patients with CODs present with physical complaints, such as insomnia, fatigue, chest pain, cardiac arrhythmia, headaches, or impotence. When other physical or psychological causes cannot be found, an SUD, an MD, or a COD should be considered. These disorders should also be considered when a patient with a chronic disease, such as chronic pain, diabetes, heart disease, gastrointestinal disorders, or hypertension, fails to respond to treatment.¹⁵

Although primary care providers should not immediately identify mental disorder symptoms as being caused by an SUD, it is important to note that many mental disorders, including mood, anxiety, sleep, and sexual disorders, can be induced by substance use. The only difference between substance-induced mental disorders and independent mental disorders is that all or most of the symptoms of a substance-induced disorder are a direct result of substance use, abuse, or withdrawal rather than mental illness. When substance-induced disorders are suspected, primary care providers should continue to evaluate psychiatric symptoms and their relationship to abstinence or ongoing substance abuse over time.¹¹

Screening

Standardized screening instruments help identify patients with potential CODs. Ideally, every patient should be screened. At the very least, any patient presenting with signs or symptoms of either a mental or a substance use problem should be evaluated for both disorders.¹⁶ Although many SUD and MD screening tools can be used in a primary care setting (e.g., the CAGE-AID,¹⁴ the Mental Health Screening Form-III,¹¹ the Primary Care Evaluation of Mental Disorders Patient Health Questionnaire^{TM20}), busy healthcare providers may find administering these tools time consuming.

Recent research has shown that one- or two-item screeners are effective in identifying those at risk for an SUD or MD (see exhibit 1).^{17, 18, 19, 20} Because the screener questions can be answered in seconds, they can be asked during routine visits. Computerized screeners, such as the Drug Abuse Problem Assessment for Primary Care^{21, 22} (DAPA-PC), are also effective when time is limited. The DAPA-PC is an Internet-based, self-administered screener for alcohol and drug use that scores patient responses, generates a patient profile for the healthcare provider, and offers motivational

Triple Threat: SUDs, MDs, and HIV/AIDS

Healthcare providers should be aware that some people with CODs are infected with HIV.²⁴ CODs may interfere with effective HIV care because of many factors, including poor adherence to antiretroviral therapy.^{24, 25} In addition, the presence of HIV infection may result in more severe co-occurring symptoms. Patients with HIV should be screened regularly for CODs. The Health Resources and Services Administration's report, *A Guide to Primary Care of People With HIV/AIDS*, provides valuable information on managing co-occurring disorders in patients with HIV/AIDS.¹³

Exhibit 1. Brief Screening Tests

Problem	Questions	Possible Responses	Positive Screen
Alcohol ¹⁸	When was the last time you had more than four (for women)/five (for men) drinks in 1 day?	1) never 2) in the past 3 months 3) over 3 months ago	in the past 3 months
Alcohol or Drugs ¹⁷	In the last year: 1) Have you ever drunk alcohol or used drugs more than you meant to? 2) Have you felt you wanted or needed to cut down on your drinking or drug use?	yes or no	yes to either question
Depression ²⁰	During the past 2 weeks: 1) Have you often been bothered by feeling down, depressed, or hopeless? 2) Have you often been bothered by little interest or pleasure in doing things?	yes or no	yes to either question

Substance Abuse in Brief Fact Sheet

Fall 2006, Volume 4, Issue 2

messages and advice to the patient. A similar screener for identifying CODs has recently been developed.²³

Treatment for Patients With COD

All patients who screen positive for a COD need a thorough assessment. The Quadrants of Care Model, a framework that classifies persons with CODs into four basic groups based on symptoms and relative symptom severity, helps determine appropriate patient care based on the type and severity of the patient's symptoms (see exhibit 2).¹¹ Patients with severe CODs or those whose symptoms worsen should be referred for further assessment and/or treatment by a specialist, but many patients may delay or refuse seeing a substance abuse treatment or mental health provider.² In addition, the gap between the availability of treatment slots and the brevity of formal treatment may result in referred patients returning to the primary care provider for COD treatment.²⁶ Some patients may need to be screened for suicidality because persons with CODs are at increased risk for committing suicide.¹¹

An effective way to address mild to moderate CODs in the primary care setting is through a brief intervention. Brief interventions—short, patient-centered interventions aimed at modifying behavior—have been shown to be effective in reducing alcohol²⁷ and drug use²⁸ as well as anxiety and depression²⁹ when used in the primary care setting. Brief interventions typically are provided over one to five visits and consist of the following:^{19, 30, 31}


1. Providing simple, concise **feedback** on patients' risk for COD based on their histories, physical and behavioral warning signs, screening test results, and information on how these disorders affect them;
2. Offering clear oral and written **advice**, including recommendations on safe alcohol consumption limits, the negative effects of alcohol and drug use, and behavior modification; and
3. Establishing a mutually consented **plan of action** that addresses SUDs and MDs as appropriate and includes specific goals for behavior change, prescription of behavioral or pharmaceutical treatments, referrals for further assessment and treatment when appropriate, and followup plans, either in person or via the telephone

The National Institute on Alcohol Abuse and Alcoholism's publication *Helping Patients Who Drink Too Much: A Clinician's Guide* and the complementary *A Pocket Guide for Alcohol Screening and Brief Intervention* provide valuable information about conducting brief interventions in the primary care setting.

Ways To Help Patients With CODs

- Provide feedback and advice in a clear, concise, nonjudgmental, and supportive manner. Empathy is particularly important because CODs are still associated with shame and guilt by many persons.³⁰
- Talk about these disorders in a matter-of-fact way—as treatable conditions—to put the patient at ease and encourage cooperation.^{30, 32}
- Become familiar with and refer patients to 12-Step or mutual-help organizations such as Alcoholics Anonymous (www.aa.org), Double Trouble in Recovery (www.doubletroubleinrecovery.org), and Narcotics Anonymous (www.na.org).
- Develop relationships with local substance abuse and mental health treatment providers, including those providing integrated treatment.
- Stay current on COD research and practices by visiting the Web sites of the Substance Abuse and Mental Health Services Administration's Co-Occurring Center for Excellence (coce.samhsa.gov), the National Mental Health Association (www.nmha.org), and the National Alliance on Mental Illness (www.nami.org), as well as by participating in COD-specific training or continuing education.

Exhibit 2. Quadrants of Care Model



Substance Abuse in Brief Fact Sheet

Fall 2006, Volume 4, Issue 2

Resources

Publications

- TIP 42: *Substance Abuse Treatment for Persons With Co-Occurring Disorders* (www.samhsa.gov)
- *Helping Patients Who Drink Too Much: A Clinician's Guide* (pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/guide.pdf)
- KAP Keys and Quick Guide for Clinicians Based on TIP 24: *A Guide to Substance Abuse Services for Primary Care Clinicians*, NCADI #s KAPT24 and QGCT24
- KAP Keys and Quick Guide for Clinicians Based on TIP 34: *Brief Interventions and Brief Therapies for Substance Abuse*, NCADI #s KAPT34 and QGCT34
- KAP Keys and Quick Guide for Clinicians Based on TIP 42: *Substance Abuse Treatment for Persons With Co-Occurring Disorders*, NCADI #s KAPT42 and QGCT42
- *HIV/AIDS: Is Your Client at Risk?* NCADI # MS965
- *Alcohol Use Among Older Adults: Pocket Screening Instruments for Health Care and Social Service Providers*, NCADI # PHD883
- *Special Feature Kit: Overcoming Addiction and Mental Disorders*, NCADI # AMDKIT
- *Building Bridges—Co-Occurring Mental Illness and Addiction: Consumers and Service Providers, Policymakers, and Researchers in Dialogue* (www.mentalhealth.samhsa.gov/publications/allpubs/SMA04-3892)
- *Improving the Quality of Health Care for Mental and Substance Use Conditions* (fermat.nap.edu/catalog/11470.html)
- *Co-Occurring Disorders: Integrated Dual-Disorders Treatment* (www.mentalhealth.samhsa.gov/cmhs/communitysupport/toolkits/cooccurring)

See ordering information for National Clearinghouse for Alcohol and Drug Information (NCADI) publications on the back cover.

Web Sites

- Substance Abuse Treatment Facility Locator (www.findtreatment.samhsa.gov; 1-800-662-HELP)
- Mental Health Services Locator (www.mentalhealth.samhsa.gov/databases)
- Co-Occurring Center for Excellence (COCE) (www.coce.samhsa.gov)
- National Mental Health Information Center (NMHIC) (www.mentalhealth.samhsa.gov)
- National Clearinghouse for Alcohol and Drug Information (NCADI) (www.ncadi.samhsa.gov)
- Substance Abuse and Mental Health Services Administration (www.samhsa.gov)

References

- ¹ Substance Abuse and Mental Health Services Administration. *Results From the 2004 National Survey on Drug Use and Health: National Findings*. NSDUH Series H-28. DHHS Publication No. (SMA) 05-4062. Rockville, MD: Office of Applied Studies, 2005.
- ² Baron, D.A., Hay, D.P., and Easom, H.M. Treating patients in primary care: The impact of mood, behavior, and thought disturbances. *Journal of the American Osteopathic Association* 103(7):319–329, 2003.
- ³ Institute of Medicine. *Improving the Quality of Health Care for Mental and Substance-Use Conditions*. Washington, DC: The National Academies Press, 2006.
- ⁴ Mertens, J.R., Lu, Y.W., Parthasarathy, S., Moore, C., and Weisner, C.M. Medical and psychiatric conditions of alcohol and drug treatment patients in an HMO: Comparison with matched controls. *Archives of Internal Medicine* 163(20):2511–2517, 2003.
- ⁵ Edlund, M.J., Unutzer, J., and Wells, K.B. Clinician screening and treatment of alcohol, drug, and mental problems in primary care: Results from healthcare for communities. *Medical Care* 42(12):1158–1166, 2004.
- ⁶ Mallin, R., Slott, K., Tumblin, M., and Hunter, M. Detection of substance use disorders in patients presenting with depression. *Substance Abuse* 23(2):115–120, 2002.
- ⁷ Kahn, L.S., Halbreich, U., Bloom, M.S., Bidani, R., Rich, E., and Hershey, C.O. Screening for mental illness in primary care clinics. *International Journal of Psychiatry in Medicine* 34(4):345–362, 2004.
- ⁸ Wilson, C.R., Sherritt, L., Gates, E., and Knight, J.R. Are clinical impressions of adolescent substance use accurate? *Pediatrics* 114(5):e536–e540, 2004.

Substance Abuse in Brief Fact Sheet

Fall 2006, Volume 4, Issue 2

- ⁹ Ferguson, L., Ries, R., and Russo, J. Barriers to identification and treatment of hazardous drinkers as assessed by urban/rural primary care doctors. *Journal of Addictive Diseases* 22(2):79–90, 2003.
- ¹⁰ Bushnell, J., McLeod, D., Dowell, A., Salmond, C., Ramage, S., Collings, S., Ellis, P., Kljakovic, M., and McBain, L. Do patients want to disclose psychological problems to GPs? *Family Practice* 22(6):631–637, 2005.
- ¹¹ Center for Substance Abuse Treatment. *Substance Abuse Treatment for Persons With Co-Occurring Disorders*. Treatment Improvement Protocol (TIP) Series 42. DHHS Publication No. (SMA) 05-3992. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2005.
- ¹² Swann, A.C., Geller, B., Post, R.M., Altshuler, L., Chang, K.D., Delbello, M.P., Reist, C., and Juster, I.A. Practical clues to early recognition of bipolar disorder: A primary care approach. *Primary Care Companion to the Journal of Clinical Psychiatry* 7(1):15–21, 2005.
- ¹³ Health Resources and Services Administration (HRSA). *A Guide to Primary Care of People With HIV/AIDS, 2004 Edition*. Rockville, MD: HRSA, 2004. hab.hrsa.gov/tools/primarycareguide [accessed December 16, 2005].
- ¹⁴ Center for Substance Abuse Treatment. *Guide to Substance Abuse Services for Primary Care Physicians*. Treatment Improvement Protocol (TIP) Series 24. DHHS Publication No. (SMA) 97-3139. Rockville, MD: Substance Abuse and Mental Health Services Administration, 1997.
- ¹⁵ Mersy, D.J. Recognition of alcohol and substance abuse. *American Family Physician* 67(7):1529–1532, 2003.
- ¹⁶ Ziedonis, D., and Brady, K. Dual diagnosis in primary care: Detecting and treating both the addiction and mental illness. *Medical Clinics of North America* 81(4):1017–1036, 1997.
- ¹⁷ Brown, R.L., Leonard, T., Saunders, L.A., and Papasouliotis, O. A two-item conjoint screen for alcohol and other drug problems. *Journal of the American Board of Family Practice* 14(2):95–106, 2001.
- ¹⁸ Canagasaby, A., and Vinson, D.C. Screening for hazardous or harmful drinking using one or two quantity-frequency questions. *Alcohol and Alcoholism* 40(3):208–213, 2005.
- ¹⁹ Fleming, M.F. Screening and brief intervention in primary care settings. *Alcohol Research & Health* 28(2):57–62, 2004/2005.
- ²⁰ Kroenke, K., Spitzer, R.L., and Williams, J.B. The Patient Health Questionnaire-2: Validity of a two-item depression screener. *Medical Care* 41(11):1284–1292, 2003.
- ²¹ Holtz, K., Landis, R., Nemes, S., and Hoffman, J. Development of a computerized screening system to identify substance abuse in primary care. *Journal for Healthcare Quality* 23(3):34–37, 45, 2001.
- ²² Nemes, S., Rao, P.A., Zeiler, C., Munly, K., Holtz, K.D., and Hoffman, J. Computerized screening of substance abuse problems in a primary care setting: Older vs. younger adults. *American Journal of Drug and Alcohol Abuse* 30(3):627–642, 2004.
- ²³ Libretto, S. “Developing and pilot testing a co-occurring disorders screening instrument for substance abuse treatment settings.” Paper presented at the American Public Health Association 133d Annual Meeting, Philadelphia, December 10–14, 2005.
- ²⁴ Douaihy, A.B., Jou, R.J., Gorske, T., and Salloum, I.M. Triple diagnosis: Dual diagnosis and HIV disease, Part 1. *The AIDS Reader* 13(7):331–341, 2003.
- ²⁵ Lucas, G.M., Gebo, K.A., Chaisson, R.E., and Moore, R.D. Longitudinal assessment of the effects of drug and alcohol abuse on HIV-1 treatment outcomes in an urban clinic. *AIDS* 14:357–366, 2002.
- ²⁶ Chychula, N.M., and Sciamanna, C. Help substance abusers attain and sustain abstinence. *The Nurse Practitioner* 27(11):30–47, 2002.
- ²⁷ Whitlock, E.P., Polen, M.R., Green, C.A., Orleans, T., Klein, J., and U.S. Preventive Services Task Force. Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: A summary of the evidence for the U.S. Preventive Services Task Force. *Annals of Internal Medicine* 140(7):557–568, 2004.
- ²⁸ Bernstein, J., Bernstein, E., Tassiopoulos, K., Heeren, T., Levenson, S., and Hingson, R. Brief motivational intervention at a clinic visit reduces cocaine and heroin use. *Drug and Alcohol Dependence* 77(1):49–59, 2005.
- ²⁹ Lang, A.J. Brief intervention for co-occurring anxiety and depression in primary care: A pilot study. *International Journal of Psychiatry in Medicine* 33(2):141–154, 2003.
- ³⁰ Finfgeld-Connett, D.L. Treatment of substance misuse in older women: Using a brief intervention model. *Journal of Gerontological Nursing* 30(8):30–37, 2004.
- ³¹ Moyer, A., and Finney, J.W. Brief interventions for alcohol problems: Factors that facilitate implementation. *Alcohol Research & Health* 28(1):44–50, 2004/2005.
- ³² National Institute on Alcohol Abuse and Alcoholism (NIAAA). *Helping Patients Who Drink Too Much: A Clinician’s Guide*. Rockville, MD: NIAAA, 2005. pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/guide.pdf [accessed December 16, 2005].

Substance Abuse in Brief Fact Sheet

Fall 2006, Volume 4, Issue 2

Substance Abuse in Brief Fact Sheet

Substance Abuse in Brief Fact Sheet is produced under contract number 270-04-7049 by JBS International, Inc., and The CDM Group, Inc., for the Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration (SAMHSA), U.S. Department of Health and Human Services (HHS). An electronic version of *Substance Abuse in Brief Fact Sheet* is available online at www.kap.samhsa.gov under Products. If you wish to reference or reproduce this issue, citation of this publication is appreciated.

Recommended Citation: Center for Substance Abuse Treatment. Identifying and Helping Patients With Co-Occurring Substance Use and Mental Disorders: A Guide for Primary Care Providers. *Substance Abuse in Brief Fact Sheet* Fall 2006, Volume 4, Issue 2.

Public Domain Notice: All material appearing in this report is in the public domain and may be reproduced or copied without permission. This publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, HHS.

DHHS Publication No. (SMA) 06-4187
Published 2006

Ordering Information

To order additional copies of *Substance Abuse in Brief Fact Sheet* and the other SAMHSA products, contact SAMHSA's NCADI
P.O. Box 2345
Rockville, MD 20847-2345
Phone: 800-729-6686, TDD: 800-487-4889
Fax: 240-221-4292
Web site: www.ncadi.samhsa.gov



National Clearinghouse for Alcohol and Drug Information
Substance Abuse and Mental Health Services Administration
P.O. Box 2345
Rockville, MD 20847-2345

PRSRT STD
U.S. POSTAGE
PAID
ODENTON, MD
PERMIT NO. 350